



ACC.18™

67th Annual Scientific Session & Expo


ORLANDO
MARCH 10 - 12
2018

A PRAGMATIC RANDOMIZED TRIAL OF CYP2C19 GENOTYPING IMPLEMENTATION FOLLOWING PERCUTANEOUS CORONARY INTERVENTION (PCI)

Sony Tuteja, PharmD, MS
Twitter @sony_tuteja

Perelman School of Medicine at the University of
Pennsylvania, Philadelphia, PA

CYP2C19 polymorphisms and clopidogrel response



CYP2C19*2 Minor allele Frequency

Europeans	0.15
African	0.15
East Asian	0.29
South Asian	0.35


Trenk D et al. Thromb Haemost 2013

Scott et al. Clin Pharm Ther 2013




ACC.18™

CYP2C19 polymorphisms and clopidogrel response



Shuldiner AR et al. JAMA 2009



Mega JL et al. JAMA 2010



ACC.18™

Conflicting recommendations exist for *CYP2C19* testing

- **Clinical Pharmacogenetics guidelines recommend prasugrel or ticagrelor in *CYP2C19* LOF carriers. ¹**
- **FDA placed a “black box” warning on the clopidogrel label in 2010 recommending alternative agents among carriers of 2 LOF alleles. ²**
- **ACC/AHA guidelines do not recommend routine *CYP2C19* genetic testing. ³**

¹ Scott SA et al. Clinical Pharmacology and Therapeutics 2013

² <https://www.fda.gov/Drugs/DrugSafety/default.htm>

³ Levine GN et al. Circulation 2016




Assessment of prospective *CYP2C19* genotype guided Dosing of AntiPlatelet Therapy in Percutaneous Coronary Intervention (ADAPT)

- **Rationale**
 - It is unknown how physicians will utilize the *CYP2C19* test results in a real world setting
- **Objective**
 - To provide evidence regarding the implementation and effectiveness of *CYP2C19* testing
- **Hypothesis**
 - Pharmacogenetic test results will influence prescribing of antiplatelet medications post PCI



The ADAPT study: *A Pragmatic Randomized Clinical Trial*



Primary endpoint

Proportion of participants receiving prasugrel/ticagrelor

Secondary endpoints

- 1. Agreement with the genotype guided antiplatelet recommendations**
- 2. Clinical Outcomes: Major Adverse Cardiac Events and Major Bleeding**



Study Intervention

- Genotyped Group
 - Buccal swab for genotyping on the SpartanRx rapid turnaround device (*CYP2C19* *2, *3, *17)
 - Genotyped guided recommendations provided verbally
 - CYP2C19* *2 or *3 carriers → prasugrel or ticagrelor
 - CYP2C19* *1 or *17 carriers → clopidogrel
 - Antiplatelet choice remained at the discretion of the treating interventional cardiologist
- Usual Care
 - Saliva collected for post study genetic analysis

Spartan **RX**
CYP2C19



| ACC.18

Inclusion/Exclusion Criteria

- **Inclusion**
 - Age ≥ 18 and ≤ 80 years
 - PCI with stent implantation
- **Exclusion**
 - Need for imminent surgery
 - History of intracranial hemorrhage, stroke
 - Active bleeding
 - Need for long-term anticoagulation
 - Study staff unavailable to conduct randomization or genotyping



Sample size determination

- Rate of pre-study prasugrel/ticagrelor use (~20%)
- Anticipated increase in prasugrel/ticagrelor prescribing based on frequency of *CYP2C19* LOF alleles (~30-35%)
- A sample size of 138 per group would provide 80% power at an $\alpha=0.05$ to detect a 15% difference in prasugrel/ticagrelor prescribing between the groups
- Sample size was increased to 250 per group to allow for subgroup comparisons



Participant Demographics

	Genotyped N=249	Usual Care N=255	P-value
Sex, n(%)			
Male	181 (73%)	190 (74%)	0.76
Race, n (%)			
White	194 (78%)	197 (77%)	0.99
Black	48 (19%)	51 (20%)	
Age	63.0 ± 9.7	62.9 ± 10.2	0.90
ACS	124 (50%)	129 (50%)	0.93



Participant History

	Genotyped N=249	Usual Care N=255	P-value
<u>Medical History, n(%)</u>			
Hypertension	190 (76)	199 (78)	0.67
Cholesterol	112 (45)	113 (44)	0.80
PCI	83 (33)	83 (33)	0.63
Diabetes	89 (35)	79 (31)	0.26
MI	63 (25)	67 (26)	0.84
CABG	32 (13)	36 (14)	0.70
<u>P2Y12 prior, n (%)</u>			
Clopidogrel	80 (32)	85 (33)	0.65
Prasugrel	12 (5)	9 (4)	
Ticagrelor	2 (1)	6 (2)	
None	154 (62)	154 (60)	



Procedure Characteristics

	Genotyped N=249	Usual Care N=255	p-value
Length of stay, mean (SD), days	2.9 ± 3.7	3.1 ± 4.0	0.66
Drug eluting stents, n(%)	237 (95)	236 (93)	0.27
Number of stents, mean (SD)	1.3 ± 0.6	1.4 ± 0.6	0.73



CYP2C19 Genotypes for Intervention Group

Genotyping results available 1.4 ± 0.4 hours post swab

Genotype	Frequency
*1/*1	34%
*1/*17	31%
*17/*17	5%
*1/*2	20%
*2/*17	5%
*2/*2	3%
Inconclusive	4%

} 28%



Primary Outcome: Antiplatelet Drugs Prescribed

	Genotyped N=249	Usual Care N=255	P-value
Clopidogrel	174 (70%)	201 (79%)	0.03
Prasugrel or Ticagrelor	75 (30%)	54 (21%)	

Fisher's exact test



Prasugrel/ ticagrelor use greater in the LOF carriers

	Genotyped		Usual Care N=255
	<u>No-LOF</u> N=174	<u>LOF carriers</u> n=68	
Clopidogrel	136 (78%)	32 (47%)	201 (79%)
Prasugrel or Ticagrelor	38 (22%)	36 (53%)	54 (21%)

P<0.001

P<0.001



ACC.18™

Agreement rate =
LOF prasugrel/ticagrelor + non-LOF clopidogrel
total number genotyped



| ACC.18™

Agreement with genotype guided recommendations

CYP2C19 diplotype	Phenotype	Clop	Pras/ Ticag	Non-agreement reasons
*1/*2 *2/*17 *2/*2	Intermediate or poor metabolizer (n=68)	32 (47%)	36 (53%)	9- Stable CAD 6- Cost 3- Contraindications 3- MD preference
*1/*1 *1/*17 *17/*17	Normal or rapid metabolizer (n=174)	136 (78%)	38 (22%)	6- Disease characteristics 6- Patient already on therapy 5- ACS 5- Recurrent events

Agreement rate 71%

Non-agreement rate 29%



ACC.18™

Prior antiplatelet therapy predicted antiplatelet drug choice independent of genotype

Prior P2Y12	OR remaining on same	95%CI	P-value
Clopidogrel	2.04	1.22, 3.45	0.007
Prasugrel/ ticagrelor	99.3	13.2, 744	<0.0001



Genotype did not influence prescribing among patient already on prasugrel or ticagrelor

Prior P2Y12	Genotype agreement rate
Clopidogrel (n=80)	76%
Prasugrel/ticagrelor (n=14)	21%
None (n=147)	73%
P-value	<0.0001



Clinical outcomes

	Genotyped (n=249)	Usual Care (n=255)	P-value
Follow-up time (months)	17.2 (7.5)	16.1 (8.2)	0.14
MACE	34 (13.7)	26 (10.2)	0.27
BARC 3 or 5 bleed	6 (2.4)	8 (3.1)	1.0

MACE= myocardial infarction, stroke, death from cardiovascular cause, stent thrombosis, urgent revascularization

BARC= Bleeding Academic Research Consortium



Summary

- ***CYP2C19* test results significantly influenced antiplatelet prescribing**
- **Genotype guided recommendations were followed 71% of the time**
- **Prior antiplatelet therapy significantly influenced the choice of antiplatelet drugs**



Conclusions

- **Antiplatelet prescribing was not universally in agreement with genotype suggested recommendations**
- **Physicians consider both clinical and genetic factors when prescribing antiplatelet agents following PCI**



Acknowledgements

ADAPT Study Team

Jay Giri

Daniel J. Rader

Henry Glick

William Matthai

Ashwin Nathan

Karen Monono

Craig Carcuffe

Karen Maslowski

Gene Chang

Taisei Kobayashi

Saif Anwaruddin

John Hirshfeld

Robert Wilensky

Howard C. Herrmann

Daniel M. Kolansky

Cardiac Cath Labs

Hospital of the University of
Pennsylvania

Penn Presbyterian Medical Center

Department of Laboratory Medicine

Funding

Penn Center for Precision Medicine



| ACC.18™

Extra slide



| ACC.18™

Post-hoc analysis

	Non-LOF (n=326)	LOF-clop (n=90)	LOF-pras/ticag (n=52)
MACE	33 (10.1)	14 (15.6)	9 (17.3)
Major bleed	6 (1.8)	4 (4.4)	1(1.9)
Non cardiac death	5 (1.5)	1 (1.1)	0 (0)
Composite	42 (12.9)	19 (21.1)	10 (19.2)

HR 1.84
95% CI 1.06 to 3.20
(p=0.03)

